

The Effect of dietary induced hypercholesterolemia on the lipid profile and some enzyme activities in Wistar male rats.

El-Khedr Mohamed Mostafa El-gamal

Department of Biochemistry, Faculty of Medicine (Dommitta), Al-Azher University Egypt

Abstract: Aim of study: To investigate the effects of high dietary soybean oil and cholesterol on the serum lipid profile, enzymes and biochemical changes in weanling Wistar rat's male. Materials and Methods: Twenty weanling male Wistar rats organised equally into two groups were used in this study. The first group of rats, were fed with controlled diet made up of the normal rat chow (C), and the second group was fed with hypercholesterolemia diet (HPC) enriched with 25% soybean oil and 1% cholesterol for eight weeks. Results: The dietary intake of the HPC diet was significantly increased the level of total cholesterol (TC), LDL-C and triglycerides (TG) in the serum of Wistar rats fed with HPC diet. gamma glutamyl transferase (GGT), aspartate amino transferase (AST) and alanine amino transferase (ALT), activities were also markedly elevated in rats fed with HPC diet; while, total protein and glucose level of rats fed with HPC diet was remarkably reduced, no significant difference was found in the high density lipoprotein cholesterol (HDL-C) and albumin contents of both groups. Conclusion: This study established that hypercholesterolemia is induced by high soybean oil/cholesterol diet, despite the fact that soybean oil has high content of polyunsaturated fatty acids. In addition, the HPC diet led to weight loss in the rats and caused injury of rats' heart and liver.

Keywords: hypercholesterolemia, enzyme activities, cholesterol, injury, soybean oil, Diet -induced, lipid profile.

1. INTRODUCTION

Hypercholesterolemia is a lipoprotein metabolic disorder characterized by high serum of LDL-C and blood (TC). It has been reported by Rerkasan et al. (2014) that hypercholesterolemia is one of the most important risk factors in the development and progression of atherosclerosis that lead to cardiovascular diseases (CVDs). Hypercholesterolemia poses a major problem to many societies as well as health professionals because of the close correlation between cardiovascular diseases and lipid abnormalities (Matos et al. 2015; Ramachandran *et al.*, 2003). Clinical trials have demonstrated that intensive reduction of plasma LDL-C levels could reverse atherosclerosis and decrease the incidence of cardiovascular diseases (CVDs) (Brown et al., 1990; Ichihashi *et al.*, 1998). Dietary factors such as continuous ingestion of high amounts of saturated fats and cholesterol are believed to be directly related to hypercholesterolemia and susceptibility to atherosclerosis (Asashina et al., 2005). Furthermore, dietary trials have revealed that the

concentration of serum cholesterol is affected by both the content and source of proteins (Seroungue et al., 1995; Forsythe, 2013).

Lipid structure, composition, configuration, in addition to excessive fat and cholesterol consumption are also believed to affect the lipid profile in the plasma (Zulet et al., 1999). Hypercholesterolemic animals are useful models for studies on cholesterol homeostasis, and drug trials to better understand the relationship between disorders in cholesterol metabolism, atherogenesis as well as possible treatments for the reduction of circulatory cholesterol levels (Pellizon, 2008; Jang and Wang, 2009). Inducing hypercholesterolemia in rats is often through a high fat, high cholesterol diet, with the fat source varying from lard to canola, coconut, soybean or palm oil. Commercial rations supplemented with cholesterol have also been used for these investigations (Doucet, 1987). The aim of the present study is to investigate the effect of the inclusion of soybean oil and cholesterol in the diet of Wistar rats. This was with the view of using our findings as a model for subsequent atherogenic studies .

2. MATERIALS AND METHODS

Animals and diets

Twenty male weanling rats weighing approximately 74.5 g each, were obtained from the Animal House at the faculty of Medicine (Dommita), University of Al-Azhar, Egypt. The animals were housed in groups of ten in a controlled environment with 12 h light and dark cycles. They were allowed free access to different dietary formulations (Table 1) and water ad libitum for eight weeks. Ten rats were fed with the control diet (C), while the second group was fed with HPC diet. The animals were weighed gravimetrically every week throughout the period of the experiment

Table 1. Composition of the control and hypercholesterolemic diets fed to the male Wistar rats

Ingredient (g/kg)	Control diet	Hypercholesterolemic diet
Casein	150.00	150.00
Corn starch	829.60	529.60
Cholesterol	0.00	15.00
Cellulose	20.00	230.00
Soybean oil	100.00	350.00
Mineral/vitamin mix	80.00	80.00

Assay kits and reagents

Kits for TC, HDL-C, LDL-C, TG, total protein, albumin, glucose, GGT, ALT and AST were obtained from Randox Laboratories Ltd., Ardmore, Co. Antrim, UK. All other reagents used were of analytical grade

Biochemical analysis

Analysis of both the control and hypercholesterolemic diets of the rats were determined using the methods described by AOAC, (2002) for crude protein, crude fat, crude fibre, and ash while carbohydrate was determined by subtracting the sum of the other nutrient parameters from 100. At the end of the eight weeks, the animals were fasted overnight, sacrificed under ether anaesthesia and blood collected from the jugular vein for analysis. The blood samples were centrifuged at $1200 \times g$ for 5 min and the serum was collected in Eppendorf tubes for analysis. The heart, liver and kidney of each animal was removed, weighed and stored at -80°C ; abdominal fat was also carefully dissected and weighed. The relative organ to body weight was calculated. TC, HDL-C and triglycerides were determined in the serum of the rats by adopting the protocol outlined in the manufacturer's assay kit from Randox Laboratories Ltd, Ardmore, Co. Antrim, UK. LDL-C was calculated using the Friedewald et al (1972) equation: $\text{LDL-C} = \text{TC} - [\text{HDL-C} + \text{TG}/5]$. Total protein, albumin, glucose and enzyme activities were also determined using assay kits and protocol from the same manufacturer

Statistical analysis

All data were expressed as mean \pm S.D and were statistically analyzed using one-way analysis of variance (ANOVA). Means were separated by the Duncan multiple test using SAS (2002). Values were considered significant at $p < 0.05$.

3. RESULTS AND DISCUSSION

Table 2 shows the proximate analysis of the control and hypercholesterolemic diets. There was no significant difference in the protein content of both diets, while there was a significant difference $P < 0.05$, in the crude fat content of the hypercholesterolemic diet when compared with the control. This was expected since the HPC diet had higher fat contents (25% soybean oil and 1% cholesterol). However, the ash and total carbohydrate content of the HPC was markedly low. It has been reported that the relative content of saturated and polyunsaturated fatty acids (PUFA) in the diet affects plasma level of cholesterol (Ohtani et al., 1990). Although metabolic studies have shown that consumption of ω -6 PUFA lowers circulating cholesterol level (Ramachandran et al., 2003). However, in this study, soybean oil, a rich source of PUFA and cholesterol without cholic acid was successfully used to induce hypercholesterolemia. There were significant differences in the weight gain pattern of the animals fed with HPC diet compared to those fed with the control diet as shown in Figure 1. While animals fed with the control diet maintained consistent weight gain throughout the period and those fed with the HPC, only gained weight up till the 4th week, then started a steady weight decline till the end of the experiment. The resultant weight loss confirmed by the organ to body weight ratio (Table 3) which could be as a result of a reduction in nutrient intake because of the high diet fat content which might impaired the absorption of protein and other nutrients (Zulet and Martinez, 1995; Woo and Henry, 1996; Matos et al., 2015). Records of similar works showed no significant changes in weight gain among the control and hypercholesterolemic animals (Ramachandran et al., 2003),

while Harnafi et al. (2009) reported a linear weight increase in all the animals fed with both experimental and control diets. However, Matos et al. (2015), and Hartvigsen et al. (2007) reported similar weight loss as observed in this study with diet-induced hypercholesterolemic rat models.

Table 2. Proximate composition of the control and hypercholesterolemic diets fed to the male Wistar rats

(Parameter (%))	Control diet	Hypercholesterolemic diet
Moisture	8.90 ± 0.14 ^b	9.15 ± 0.07 ^a
Ash	10.24 ± 0.16 ^a	7.73 ± 0.04 ^b
Crude protein	25.95 ± 0.14 ^a	27.74 ± 0.02 ^a
Crude fat	18.64 ± 0.04 ^b	32.44 ± 0.14 ^a
Crude fiber	3.99 ± 0.14 ^b	4.25 ± 0.14 ^a
Total carbohydrate	59.53 ± 0.22 ^a	47.25 ± 0.13 ^b

Values are mean ± SD of 3 replicates. a - b Test values along the same row carrying different superscripts for each parameter are significantly different (P < 0.05)




Figure 1. Growth response of male Wistar rats fed with control and hypercholesterolemic (HPC) diet.

However, Matos et al. (2015), and Hartvigsen et al. (2007) reported similar weight loss as observed in this study with diet-induced hypercholesterolemic rat models. The marked reduction in the abdominal fat observed in the HPC fed rats could be as a result of ketogenic mechanisms (Zulet et al., 1999). Table 4 shows that there was a significant elevation of TC, LDLC and triglycerides in the HPC fed rats, while there was no such difference in the HDL-C of both groups. Increase of LDL-C level is considered as one of the risk factors for the development of atherosclerosis and related cardiovascular diseases (Getz and Reardon, 2006). Hence, the result obtained in this study indicated that the dietetic model is suitable for atherogenesis studies. High serum triglyceride levels has also been reported to be an important risk factor as it influences lipid deposition and clotting mechanisms (Harnafi et al., 2009). Many studies have reported high dietary fat

and cholesterol induce hypercholesterolemia in animal models (Cherng and Shih, 2005; Martinello et al., 2006). Similar results were also observed with the HPC fed animals having elevated lipid status than the control. This was an indication that hypercholesterolemia was successfully established in this study.

Table 3. Effect of hypercholesterolemic diet on organ to body weight of male Wistar rats.

Organ (g)	Control rats	Hypercholesterolemic rats
Body weight	167.84 ± 4.01 ^a	102.50 ± 12.50 ^b
Liver	10.00 ± 1.41 ^a	8.18 ± 0.25 ^b
Heart	3.53 ± 0.18 ^a	2.89 ± 0.09 ^b
Abdominal fat	5.97 ± 0.69 ^a	3.80 ± 0.08 ^b
Kidney	3.00 ± 0.00 ^a	3.00 ± 0.00 ^a

Values are mean ± SD of 3 replicates. a - b Test values along the same row carrying different superscripts for each parameter are significantly different (P < 0.05)

Table 4. Effect of hypercholesterolemic diet on serum lipid profile of male Wistar rats

Parameters (mmol/l)	Control rats	Hypercholesterolemic rats
TC	2.50 ± 0.57 ^b	5.10 ± 0.28 ^a
HDL-C	0.49 ± 0.21 ^a	0.49 ± 0.28 ^a
LDL-C	2.00 ± 0.35 ^b	4.50 ± 0.28 ^a
TG	1.30 ± 0.28 ^b	3.50 ± 0.28 ^a

Values are mean ± SD of 3 replicates. a-b Test values along the same row carrying different superscripts for each parameter are significantly different (P < 0.05) .

Table 5. Effect of Hypercholesterolemic diet on liver and heart functions of male Wistar rats

Parameter	Control rats	Hypercholesterolemic rats
Total protein (g/l)	73.50 ± 1.41 ^a	62.00 ± 1.41 ^b
Albumin (g/l)	42.50 ± 1.41 ^a	42.00 ± 1.41 ^a
Glucose (mmol/l)	7.18 ± 0.39 ^a	4.15 ± 0.21 ^b
GGT (U/l)	19.50 ± 0.00 ^b	38.95 ± 0.18 ^a
AST (U/l)	40.25 ± 0.35 ^b	47.25 ± 0.35 ^a
ALT (U/l)	9.50 ± 0.35 ^b	27.50 ± 0.71 ^a

Values are mean \pm SD of 3 replicates. a - b Test values along the same row carrying different superscripts for each parameter are significantly different ($P < 0.05$).

The data in Table 5 indicates that there was no significant difference ($P < 0.05$) in the albumin level of both the animals fed on the control diet and those fed on the hypercholesterolemic diet. Low serum protein was observed in the HPC fed rats and could be due to poor feed intake and utilization. The high fat diet, because of the energy density, might have led to early satiety, reduced feed intake and impaired absorption of protein and other nutrients (Rolls, 2000). The major reason for the marked decrease ($P < 0.05$) in serum glucose in the HPC fed rats was uncertain but it could be that the high serum cholesterol increases the level of glucagon-like peptide-1 which enhances insulin secretion from the pancreatic beta-cells leading to hypoglycemia (Xu et al., 1999; Prasad, 2008). It could also be as a result of reduced supply of glucose through the diet and subsequent demand for pentose and glycogen formation.

Reports have indicated that saturated fats and polyunsaturated fats may work together to regulate the expression of several enzymes involved in carbohydrate metabolism (Jump et al., 1994). The enzyme activities shown in Table 5 indicated that all three enzymes (GGT, AST and ALT) were elevated in the serum of the HPC fed rats. This could be as a result of leakage of the enzymes into the serum as a result of damage heart and liver tissue damage. Elevated serum activity of these enzymes has been reported to be an indicators of calculated risk of hepatocellular and cardiovascular disease. According to Pincus and Schaffner (1996), AST and ALT are released into serum when there is severe hepatocellular injury. Elevated serum ALT levels in the absence of viral hepatitis and alcoholism has been reported to lead to a higher risk of cardiovascular disease (Ioannou, 2006). The heart also has a high AST content which becomes elevated in myocardial infarction. These reports agree with the present study which showed hepatic injury and cardiovascular distress in the rats fed with HPC. Ruttman et al. (2005), reported a correlation between GGT and cardiovascular mortality indicating that the higher the elevation of GGT, the greater the risk of death .

4. CONCLUSION

In This study, our results has revealed that dietary intake of hypercholesterolemia diet (HPC) enriched with 25% soybean oil and 1% cholesterol as a source of fat induced hypercholesterolemia despite of its high content of both mono and polyunsaturated fatty acids. Hypercholesterolemia led to reduced serum glucose level with no significant effect on serum albumin and HDL-C induce weight loss and malnutrition in the rats model. Also, the dietary intake of soybean oil and cholesterol led to impaired hepatic and cardiovascular distress in the rats fed with the hypercholesterolemic diet

REFERENCES

1. AOAC (2002). Official Methods of Analysis. Association of Analytical Chemists. 17th edn, Washington DC.
2. Asashina M, Sato M, Imaizumi K (2005). Genetic analysis of diet induced hypercholesterolemia in exogenously hypercholesterolemic (ExHC) rats. *J. Lipid. Res.*, 46: 2289-2294. doi:10.1194/jlr.M500257JLR200.
3. Brown AJ, Jessup W (1999). Oxysterols and atherosclerosis. *Atherosclerosis*. 142: 1-28.
4. Cherng JY, Shih MF (2005). Preventing dyslipidemia by *chorella pyrenioidosa* in rats and hamsters after chronic high fat diet treatment. *Life Sci.*, 76: 3001-13.
5. Doucet C, Flament C, Sautier C, Lemonnies D (1987). Effect of an hypercholesterolemic diet on the level of several serum lipids and apolipoproteins in nine rat strains. *Reprod. Nutr. Dev.* 27: 897-906.
6. Forsythe WA (2013). Soy protein, thyroid regulation and cholesterol metabolism. *J. Nutr.*, 125: 619-623.
7. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499-502.
8. Getz GS, Reardon CA (2006). Diet and murine atherosclerosis. *Arteriosclerosis, Thrombosis Vascular Biol.*, 26: 242-249 doi:10.1161/01.ATV.0000201071.49029.17.
9. Harnafi H, Aziz M, Amrani S (2009). Sweet Basil (*Ocimum basilicum* L.) improves lipid metabolism in hypercholesterolemic rats. *E- SPEN, the European e-J. Clinical Nutr. Metabolism* 4:e181-e186 doi:10.1016/j.eclnm.2009.05.011.
10. Hartvigsen K, Binde CJ, Hansen LF, Rafia A, Juliano J, Horko S, Steinberg D, Palinski W, Witztum JL, Li AC (2007). A diet-induced hypercholesterolemic murine model to study atherogenesis without obesity and metabolic syndrome. *Arterioscler Thromb Vasc. Biol.*, 27: 878-885 doi: 10.1161/01.ATV.0000258790.3581002.
11. Ichihashi T, Izawa M, Miyata K, Mizui T, Hirano K, Takagishi Y (1998). Mechanisms of hypercholesterolemic action of S-89211 in rats: S8921 inhibits ileal bile acid absorption. *J. Pharmacol. Exper. Therapeutics, (JPET)* 284(1): 43-50.
12. Ioannou GN, Weiss NS, Boyko EJ, Mozaffarian D, Lee SP (2006). Elevated serum Alanine aminotransferase activity and calculated risk of coronary Heart Disease in the United States. *Hepatology*, 43(5): 1145-1151.
13. Jang MJ, Wang MH (2009). Effect of fermented soybean-derived chung koo kjang on diet-induced hyperlipidemia in Bio FIB hamsters. *Food Biotechnol.*, 23(1): 74-82. doi:10.1080/08905430802672020.
14. Jump DB, Clarke SD, Thelen A, LiiMatta M (1994). Coordinate regulation of glycolytic and lipogenic gene expression by polyunsaturated fatty acids. *J. Lipid Res.*, 35: 1076-1084.

15. Martinello F, Soares SM, Franco JJ, Santos AC, Sugohara A, Garcia SB, Curtis, Uyemura SA (2006). Hypolipemic and antioxidant activities from *Tamarindus indica* L; pulp fruit extract in hypercholesterolemic hamsters. *Food Chem. Toxicol.* 44(6): 810818.
16. Matos SL, Paula H, Pedrosa ML, Santos RC, Oliveira EL, Chianca Jr DA, Silva ME (2015). Dietary models for inducing hypercholesterolemia in rats. *Brazilian Archives Biol. Technol.*, 48(2): 203-209.
17. Ohtani H, Hayashi K, Hirata Y, Dojo S, Nakashima K, Nishio E, Kurushima H, Saeki M, Kajiyama G (1990). Effects of dietary cholesterol and fatty acids on plasma cholesterol level and hepatic lipoprotein metabolism. *J. Lipid Res.*, 31: 1413-1422.
18. Pellizon MA (2008). Diet-induced atherosclerosis/hypercholesterolemia in rodent models. Brief scientific literature review. Retrieved from www.researchdiets.com/pdf/product/atherosclerosis_3%page.pdf.
19. Pincus MR, Schaffner JR (1996). Assessment of liver function in clinical diagnosis and management by laboratory methods. Philadelphia: W.B. Saunders Company.
20. Prasad K (2008). Serum biochemical changes in rabbits on a regular diet with and without flax lignin complex following a high-cholesterol diet. *Int. J. Angiol.*, 17(1): 27-32.
21. Ramachandran HD, Narasimhamurthy K, Raina PL (2003). Modulation of cholesterol induced hypercholesterolemia through dietary factors in Indian desert gerbils (*Meriones hurricinae*). *Nutr. Res.*, 23: 245-256.
22. Rerkasem K, Gallagher PJ, Grimble RF, Calder PC, Shearman CP (2014). Managing hypercholesterolemia and its correlation with carotid plaque morphology in patients undergoing carotid endarterectomy (A Review). *Vascular Health Risk Manage.*, 4(6):1259-1264.
23. Rolls BJ (2000). The role of energy density in the over consumption of fat. *J. Nutr.*, 130: 268S-271S.
24. Ruttman E, Brant LJ, Concin H, Diem G, Rapp K, Ulmer H (2005). Gamma glutamyl transferase as a risk factor for cardiovascular mortality: An epidemiologic investigation in a Cohort of 163,944 Austrian adults. *Circulation*, 112: 2130-2137.
25. SAS (2002). SAS Systems for windows 9.0, SAS INSTITUTE INC. CARY, NC, USA.
26. Serouge C, Felgines C, Ferezon J, Hajri T, Bertin C, Mazur A (1995). Hypercholesterolemia induced by cholesterol or cysteine-enriched diets is characterized by different plasma lipoprotein and apoprotein concentrations in rats. *J. Nutr.*, 125: 619-623.
27. Woo J, Henry JB (1996). Metabolic intermediates and inorganic ions in clinical diagnosis and management by laboratory animals. Philadelphia: W.B Saunders Company, pp. 162-193.
28. Xu G, Stoffers DA, Habener JF, Bunner-Weir S (1999). Exendin-4 stimulates both beta-cell replication and neogenesis, resulting in increased beta-cell mass and improved glucose tolerance in diabetic rats. *Diabetes*, 48: 2270-2276.

29. Zulet MA, Barber A, Garcin H, Higuera P, Martinez JA (1999). Alterations in carbohydrate and lipid metabolism induced by a diet rich in coconut oil and cholesterol in a rat model. *J. AM. Coll. Nutr.*, 18(1): 36-42. -
30. Zulet MA, Martinez JA (1995). Corrective role of chickpea intake on a dietary-induced model of hypercholesterolemia. *Plant Foods Human Nutr.*, 48: 269-277.